

## Chapter 10: Gravitation

NCERT Exemplar · Class 9 · Science

---

### Multiple Choice Questions

#### Question 1

Two objects of different masses falling freely near the surface of moon would

- (a) have same velocities at any instant
- (b) have different accelerations
- (c) experience forces of same magnitude
- (d) undergo a change in their inertia

**Soln:**

Answer is (a) have same velocities at any instant

**Explanation:**

Acceleration of an object depends on acceleration due to gravity irrespective of its mass. Hence object under free fall have the same velocities.

2. The value of acceleration due to gravity

- (a) is same on equator and poles
- (b) is least on poles
- (c) is least on equator
- (d) increases from pole to equator

**Soln:**

Answer is (c) is least on equator

**Explanation:**

The value of acceleration due to gravity is least on equator because distance between surface of the earth and its centre is more on equator than in poles.

3. The gravitational force between two objects is  $F$ . If masses of both objects are halved without changing distance between them, then the gravitational force would become

- (a)  $F/4$
- (b)  $F/2$
- (c)  $F$
- (d)  $2F$

**Soln:**

Answer is (a)  $F/4$

4. A boy is whirling a stone tied with a string in an horizontal circular path. If the string breaks, the stone

- (a) will continue to move in the circular path
- (b) will move along a straight line towards the centre of the circular path
- (c) will move along a straight line tangential to the circular path
- (d) will move along a straight line perpendicular to the circular path away from the boy

**Soln:**

Answer is (c) will move along a straight line tangential to the circular path

**Explanation:**

At any instance of time object in circular motion tend to be in rectilinear motion. Object keeps on moving due to centripetal force and it moves along a straight line tangential to the circular path when strings breaks.

5. An object is put one by one in three liquids having different densities. The object floats with  $\frac{12}{3}$ , and  $\frac{9}{11}$  parts of their volumes outside the liquid surface in liquids of densities  $d_1$ ,  $d_2$  and  $d_3$  respectively. Which of the following statement is correct?

- (a)  $d_1 > d_2 > d_3$
- (b)  $d_1 > d_2 < d_3$
- (c)  $d_1 < d_2 > d_3$
- (d)  $d_1 < d_2 < d_3$

**Soln:**

Answer is (d)  $d_1 < d_2 < d_3$

6. In the relation  $F = G \frac{M m}{d^2}$ , the quantity  $G$

- (a) depends on the value of  $g$  at the place of observation
- (b) is used only when the earth is one of the two masses
- (c) is greatest at the surface of the earth
- (d) is universal constant of nature

**Soln:**

Answer is (d) is universal constant of nature

**Explanation:**

$G$  is called as Newton's constant. It is the force of gravity on a body. Value of  $G$  is  $6.66 \times 10^{-11} \text{ Nm}^2\text{kg}^{-2}$

7. Law of gravitation gives the gravitational force between

- (a) the earth and a point mass only
- (b) the earth and Sun only
- (c) any two bodies having some mass
- (d) two charged bodies only

**Soln:**

Answer is (c) any two bodies having some mass

8. The value of quantity  $G$  in the law of gravitation

- (a) depends on mass of earth only
- (b) depends on radius of earth only
- (c) depends on both mass and radius of earth
- (d) is independent of mass and radius of the earth

**Soln:**

Answer is (d) is independent of mass and radius of the earth

**Explanation:**

$G$  is an universal constant hence it is independent of mass and radius of the earth.

9. Two particles are placed at some distance. If the mass of each of the two particles is doubled, keeping the distance between them unchanged, the value of gravitational force between them will be

(a) 14 times

(b) 4 times

(c) 12 times

(d) unchanged

**Soln:**

**Answer is (b) 4 times**

**10. The atmosphere is held to the earth by**

(a) gravity

(b) wind

(c) clouds

(d) earth's magnetic field

**Soln:**

**Answer is (a) gravity**

**11. The force of attraction between two unit point masses separated by a unit distance is called**

(a) gravitational potential

(b) acceleration due to gravity

(c) gravitational field

(d) universal gravitational constant

**Soln:**

**Answer is (d) universal gravitational constant**

**Explanation:**

$$F = G \times$$

$$\frac{m_1 \times m_2}{r^2}$$

**Here point masses are separated by unit distance**

**Hence  $m_1$ ,  $m_2$  and  $r = 1$**

**Hence  $F = G$  which is a universal constant hence answer is universal gravitational constant**

**12. The weight of an object at the centre of the earth of radius  $R$  is**

- (a) zero
- (b) infinite
- (c) R times the weight at the surface of the earth
- (d)  $1/R^2$  times the weight at surface of the earth

**Soln:**

**Answer is (a) zero**

**Explanation:**

At the centre of the earth acceleration due to gravity is zero. Since weight is the product of mass and gravity. Weight of the object at the centre of the earth will be zero.

**13. An object weighs 10 N in air. When immersed fully in water, it weighs only 8 N. The weight of the liquid displaced by the object will be**

- (a) 2 N
- (b) 8 N
- (c) 10 N
- (d) 12 N

**Soln:**

**Answer is (a) 2 N**

**Explanation:**

**Object of weight displaced by liquid = weight in air - weight in liquid**

$$= 10\text{N} - 8\text{N}$$

$$= 2\text{N}$$

**14. A girl stands on a box having 60 cm length, 40 cm breadth and 20 cm width in three ways. In which of the following cases, pressure exerted by the brick will be**

- (a) maximum when length and breadth form the base
- (b) maximum when breadth and width form the base
- (c) maximum when width and length form the base
- (d) the same in all the above three cases

**Soln:**

Answer is (b) maximum when breadth and width form the base

Explanation:

Surface area and pressure are inversely proportional. Pressure will be maximum when surface area is minimum hence the answer is maximum when breadth and width form the base.

15. An apple falls from a tree because of gravitational attraction between the earth and apple. If  $F_1$  is the magnitude of force exerted by the earth on the apple and  $F_2$  is the magnitude of force exerted by apple on earth, then

- (a)  $F_1$  is very much greater than  $F_2$
- (b)  $F_2$  is very much greater than  $F_1$
- (c)  $F_1$  is only a little greater than  $F_2$
- (d)  $F_1$  and  $F_2$  are equal

Soln:

Answer is (d)  $F_1$  and  $F_2$  are equal

Explanation:

Newton's third law of motion states that for every action there is equal and opposite reaction. Hence  $F_1$  and  $F_2$  are equal.

### Solution

Solution coming soon.

---

## Short Answer Questions

### Question 16

What is the source of centripetal force that a planet requires to revolve around the Sun? On what factors does that force depend?

Soln:

Gravitational force is the source of centripetal force required to revolve around the sun. This force depends on the distance between the planet and sun along with their masses. If this force becomes zero as a result of the absence of centripetal force, the planet would shift to moving tangentially outwards to the circular route.

17. On the earth, a stone is thrown from a height in a direction parallel to the earth's surface while another stone is simultaneously dropped from the same height. Which stone would reach the ground first and why?

**Soln:**

Both the stones reach the ground simultaneously as they are dropped from the same height and their initial velocity will be the same.

18. Suppose gravity of earth suddenly becomes zero, then in which direction will the moon begin to move if no other celestial body affects it?

**Soln:**

If there is no gravitational pull from the earth, the moon starts to move in a straight line tangent to its circular path.

19. Identical packets are dropped from two aeroplanes, one above the equator and the other above the north pole, both at height  $h$ . Assuming all conditions are identical, will those packets take same time to reach the surface of earth. Justify your answer.

**Soln:**

The value of 'g' – acceleration due to gravity is constant, but depending upon the surface of the earth it varies from place to place as earth is not completely spherical. As it is flattened at the poles, the value of 'g' is maximum at the poles and the bulging at the equator causes the 'g' value to be minimum at the equator. The 'g' value increases as we move towards the poles. Hence, the packets falls gradually at the equator in comparison to the poles. Thereby, the packets stay in air for longer when dropped at the equator.

20. The weight of any person on the moon is about  $1/6$  times that on the earth. He can lift a mass of 15 kg on the earth. What will be the maximum mass, which can be lifted by the same force applied by the person on the moon?

**Soln:**

Weight of person on moon =  $1/6^{\text{th}}$  of weight on earth

Therefore, 'g' on moon =  $1/6^{\text{th}}$  'g' on earth

The force that is applied by the man to lift mass 'm' is

$$F = mg = 15g \quad (\text{on earth})$$

If he can lift a certain mass 'm' by applying the same force on moon, then

$$F = 15 \times 6 = 90\text{kg}$$

This proves acceleration due to gravity on the moon is  $1/6^{\text{th}}$  of acceleration due to gravity on earth. Hence the person can lift a mass 6 times heavier on moon than on earth.

21. Calculate the average density of the earth in terms of  $g$ ,  $G$  and  $R$ .

Soln:

$$g = GM/R^2$$

$$M = gR^2/G$$

Density of earth  $D = \text{mass/volume}$

$$= Gr^2/G \times V_e$$

$$= gR^2/G4\pi R^3$$

$$= 3g/4\pi GR$$

22. The earth is acted upon by gravitation of Sun, even though it does not fall into the Sun. Why?

Soln:

Sun provides enough centripetal force to keep the earth in its orbit and the earth provides centrifugal force due to its motion. These two forces balance each other which prevents the earth from falling into the sun.

### Solution

Solution coming soon.

## Long Answer Questions

### Question 23

How does the weight of an object vary with respect to mass and radius of the earth. In a hypothetical case, if the diameter of the earth becomes half of its present value and its mass becomes four times of its present value, then how would the weight of any object on the surface of the earth be affected?

Soln:

Let  $R$  and  $M$  be the radius and mass of the earth

Then

$$\text{Weight of the object} = M/R^2$$

Original weight  $W_0 = mg = \frac{GMm}{R^2}$

Hypothetically  $4m$  and  $R$  becomes  $R/2$

Then, weight =  $\frac{GM(4m)}{(R/2)^2}$

$= (16m g) \frac{M}{R^2}$

$= 16W_0$

Weight will be 16 times.

24. How does the force of attraction between the two bodies depend upon their masses and distance between them? A student thought that two bricks tied together would fall faster than a single one under the action of gravity. Do you agree with his hypothesis or not? Comment.

Soln:

The hypothesis is incorrect. Force of attraction between two masses separated by distance  $r$  is given by Newton law of gravitation where  $F = \frac{Gm_1m_2}{R^2}$  Where  $G$  is gravitational force and it is an universal constant.

Gravitational force is directly proportional to the product of the masses of two bodies and inversely proportional to the square of the distance between them.

Bodies thrown from same height fall with same speed irrespective of their mass. This is due to acceleration due to gravity.

$g = \frac{GM}{R^2}$  Where  $M$  is mass  $R$  is radius of the earth

This equation shows that acceleration due to gravity depends on the mass of the earth and radius of the earth. Hence tow brick tied will not fall faster than the single one.

25. Two objects of masses  $m_1$  and  $m_2$  having the same size are dropped simultaneously from heights  $h_1$  and  $h_2$  respectively. Find out the ratio of time they would take in reaching the ground. Will this ratio remain the same if (i) one of the objects is hollow and the other one is solid and (ii) both of them are hollow, size remaining the same in each case. Give reason.

Soln:

$h_1 =$

$$\frac{1}{2}gt^2$$

$h_2 =$

$$\frac{1}{2}gt^2$$

2

as  $x = 0$ 

$$\frac{t_1}{t_2}$$

2

As the acceleration remains the same, the ratio between two objects remains the same. In this case, acceleration does not depend upon mass and size.

26. (a) A cube of side 5 cm is immersed in water and then in saturated salt solution. In which case will it experience a greater buoyant force. If each side of the cube is reduced to 4 cm and then immersed in water, what will be the effect on the buoyant force experienced by the cube as compared to the first case for water. Give reason for each case.

(b) A ball weighing 4 kg of density  $4000 \text{ kg m}^{-3}$  is completely immersed in water of density  $1000 \text{ kg m}^{-3}$ . Find the force of buoyancy on it. (Given  $g = 10 \text{ m s}^{-2}$ )

Soln:

(i) Buoyant force,  $F = V\rho g$ 

$\rho$  = Density of water,  $V$  = Volume of water displaced by the body

Volume and density of an object decides its Buoyancy. Cube will experience a greater buoyancy in a saturated solution. If the cube is reduced to 4 cm on each side, the volume of cube becomes less as the buoyancy will be reduced as buoyant force is directly proportional to volume.

(ii) The magnitude of the buoyant force given by  $F = V\rho g$ 

where  $V$  = Volume of body immersed in water or volume of water displaced,  $\rho$  = Density of liquid.

-3

Hence Volume of solid =

$$\frac{\text{mass}}{\text{density}}$$

=

$$\frac{4}{4000}$$

=

$$\frac{1}{1000}$$

Buoyancy  $F = V\rho g$ 

=

$$\frac{1}{1000}$$

= 10N

## Solution

Solution coming soon.

---